(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 20 December 2001 (20.12.2001)

PCT

(10) International Publication Number WO 01/95944 A2

(51) International Patent Classification⁷: A61K 47/48

(21) International Application Number: PCT/US01/18869

(22) International Filing Date: 12 June 2001 (12.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/211,036 12 June 2000 (12.06.2000) US

(71) Applicant and

(72) Inventor: MILLS, Randell, L. [US/US]; 493 Old Trenton Road, Cranbury, NJ 08512-5601 (US).

(74) Agents: DECONTI, Giulio, A., Jr. et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

 without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



11/95944 A2

(54) Title: PRO DRUGS FOR SELECTIVE DRUG DELIVERY

PRO DRUGS FOR SELECTIVE DRUG DELIVERY

RELATED APPLICATIONS

This application claims the benefit under 35 USC 119(e) of prior-filed provisional patent application No. 60/211,036, filed June 12, 2000, entitled "Pro Drugs for Selective Drug Delivery". This application is also related to U.S. Patent Nos. 5,428,163 and 5,773,592. The entire content of the above-referenced applications are incorporated herein by this reference.

5

10

15

20

25

30

BACKGROUND OF THE INVENTION

The effects of the preponderance of drugs result from their interaction with functional macromolecular components of the organism. Such interaction alters the function of the pertinent cellular component and thereby initiates the series of biochemical and physiological changes that are characteristic of the response to the drug. The term "receptor" denotes the component of the organism with which the chemical agent interacts. There are fundamental corollaries to the statement that the receptor for a drug can be any functional macromolecular component of the organism. One is that a drug is potentially capable of altering the rate at which any bodily function proceeds; a second is that, by virtue of interactions with specific receptors, drugs do not create effects but merely modulate the rates of ongoing functions. A simple pharmacological dictum thus states that a drug cannot impart a new function to a cell. Functional changes due to a drug result from either enhancement or inhibition of the unperturbed rate. Furthermore, a drug that has no direct action can cause a functional change by competition for a binding site with another, active regulatory ligand of the receptor. Drugs are termed agonists when they cause effects as a result of direct alteration of the fundamental properties of the receptor with which they interact. Compounds that are themselves devoid of intrinsic pharmacological activity, but cause effects by inhibition of the action of a specific agonist (e.g. by competition for agonist binding sites) are designated as antagonists.

At least from a numerical standpoint, the proteins of the cell form the most important class of drug receptors. Examples include the enzymes of crucial metabolic or regulatory pathways (e.g., tyrosine hydroxylase; 3-hydroxy-3-methylglutaryl-CoA reductase). Of equal interest are proteins involved in transport processes (e.g. Ca²⁺ - ATPase; Na⁺-K⁺-ATPase) or those that are protein kinases which activate other proteins as a consequence of their binding a secondary messenger such as cAMP. Specific

5

10

15

20

25

30

binding properties of other cellular constituents can be exploited. Thus, nucleic acids are important drug receptors, particularly for chemotherapeutic approaches to the control of malignancy, and plant lectins shown remarkable specificity for recognition of specific carbohydrate residues in polysaccharides and glycoproteins. Small ions such as Ca²⁺ which can function as a regulatory ion or Fe²⁺ which can serve as an essential enzymatic cofactor can be exploited as drug receptors. And, drugs can also produce a functional change by a nonreceptor-mediated action. Certain drugs that are structural analogues of normal biological constituents may be incorporated into cellular components and thereby alter their function. This has been termed a "counterfeit incorporation mechanism" and has been implemented with analogues of purines and pyrimidines that can be incorporated into nucleic acids and that have utility in cancer chemotherapy and that have antiviral activity. Also, specific constituents of pathogens can be exploited as receptors. For example, the electron carriers of bacteria can serve as receptors as described U.S. Application No. 948,326, which is incorporated herein by reference, and the replicative enzymes of viruses can be serve as receptors for the virus HIV. Many compounds are known which have receptor or nonreceptor mediated in vitro activity as appears in The Handbook of Enzyme Inhibitors, Mahendra Kumor Jain, 1982, Wiley Interscience, New York, incorporated herein by reference. However, only a small percentage produce the desired functional change in vivo or have a high therapeutic ratio, because they are toxic in their free form; they are rapidly inactivated or excreted; or, they cannot obtain access to their target receptor or site of action because they are impermeant to cells or biological barriers such as the blood brain barrier due to unfavorable energetics due, for example, to the possession of polar or charge groups; or, they are toxic as a consequence of being nonselective with regards to their access to and action with receptors in one biological environment or compartment relative to another. In these cases, compounds which demonstrate in vitro efficacy are ineffective therapeutics.

SUMMARY OF THE INVENTION

The present disclosure relates to chemical compositions and methods for delivering biologically active agents to a biological compartment. In an embodiment, the invention relates to a chemical compound having the formula A-B, where A is a moiety capable of receiving energy and B is a biologically active agent covalently

5

10

15

20

25

30

bonded to B, wherein the bond between A and B is capable of heterolytic cleavage upon reception of energy by A.

In an advantageous embodiment, the energy causes A to achieve an excited state, the relaxation of which causes heterolytic cleavage of the bond between A and B. In another embodiment, A is a photochromic or thermochromic moiety. In yet another embodiment, the chemical compound is 1-phosphonoformate,1,5-di-(p-N-ethyl-N-ethylaminophenyl)-1,5-bis-(p-N,N-dimethylaniline)-1,3-pentadiene.

The invention also pertains to a method for selectively delivering a biologically active agent to a biological compartment, which comprises introducing a chemical compound having the formula A-B, where A is a carrier moiety and B is the biologically active agent, into a biological compartment and exposing the chemical compound to an energy sufficient to cause heterolytic cleavage of the bond between A and B, thereby releasing B from A.

DETAILED DESCRIPTION OF THE INVENTION

The invention relates to a chemical compound having the formula A-B, where A is a moiety capable of receiving energy and B is a biologically active agent covalently bonded to B, wherein the bond between A and B is capable of heterolytic cleavage upon reception of energy by A.

In a preferred embodiment, the chemical compound is a prodrug, e.g., that permeates the desired biological compartment and undergoes release of a biologically active agent, in an active form, inside of the desired compartment. In a further embodiment, the prodrug achieves a greater therapeutic effect or therapeutic ratio relative to the free drug alone as a consequence of altered pharmacokinetics or pharmacodynamics (such as a desirable kinetics of release, a resistance to inactivation or excretion, greater solubility, enhanced absorption, a diminished toxicity, or greater access to the cellular or biological compartment which is the site of action of the drug).

In an advantageous embodiment, A is a photochromic or thermochromic molecule, and B is a biologically active agent such as a drug moiety. In yet another embodiment, A comprises a cationic dye which demonstrates photochromic behavior with electromagnetic radiation and bleaching agents. Some examples of cationic dyes which can be used for A are di or triarylmethane dyes, triarylmethane lactone or cyclic ether dyes, cationic indoles, pyronines, phthaleins, oxazines, thiazines, acridines,

5

10

15

20

25

30

phenazines, anthocyanidins, cationic polymethine dyes, azo or diazopolymethines, styryls, cyanines, hemicyanines, and dialkylaminopolyenes.

In a preferred embodiment, the chemical compound is a luminide. A luminide comprises a universal carrier molecule linked to one or more of virtually any biologically active agent including known pharmaceuticals and pesticides. The luminide conjugate potentiates delivery to the desired biological compartment and potentiates intracellular uptake of the biologically active agent which breaks apart due to a reversible bond between the carrier moiety and the biologically active agent to result in release of the free biologically active agent, allowing the desired pharmacological effect to occur. An embodiment of a luminide is a cellular permeant prodrug where intracellular drug release occurs when the prodrug undergoes heterolytic cleavage of the bond between a drug moiety and a carrier moiety. In another embodiment, the luminide is a two-part molecule where each part is a functionality with a defined purpose.

An exemplary luminide is of the structure A-B, where A represents a functionality which forms a reversible bond with B, which is released through heterolytic cleavage of the covalent bond of A with B. B is, in an embodiment, a drug moiety which is released in its free form into the environment. The free drug moiety effects a therapeutic functional change in the system to which it is introduced. Such mechanisms may include receptor mediated mechanisms including reversible and irreversible competitive agonism or antagonism, e.g., a molecule known as a "suicide substrate", a transition state analogue, or a noncompetitive or uncompetitive agonism or antagonism, or the mechanism may be a nonreceptor mediated mechanism such as a "counterfeit incorporation mechanism". The heterolytic cleavage releases the drug moiety into the desired compartment in active form to effect a greater therapeutic effect or therapeutic ratio relative to the free drug moiety alone as a consequence of altered pharmacokinetics or pharmacodynamics such as a desirable kinetics of release, a resistance to inactivation or excretion, greater solubility, enhanced absorption, a diminished toxicity, or greater access to the cellular or biological compartment which is the site of action of the drug moiety. In an embodiment, the B moiety is a bleaching agent (a molecule which covalently bonds to A, the carrier moiety of the Luminide such as a photochromic functionality). Such agents comprise essentially any nucleophilic group including phosphate, sulfide, sulfite, sulfate, carboxylate, hydroxyl, or amine.

5

10

15

20

25

30

The invention also pertains to a method for selectively delivering a biologically active agent to a biological compartment, which comprises introducing a chemical compound having the formula A-B, where A is a carrier moiety and B is the biologically active agent, into a biological compartment and exposing the chemical compound to an energy sufficient to cause heterolytic cleavage of the bond between A and B, thereby releasing B from A.

In another embodiment, effective drug delivery is achieved based on the stability of the A-B reversible bond which has a sufficient stability (i.e., a long enough half-life) to permit the prodrug to penetrate the desired biological compartment before the drug moiety is released through heterolytic cleavage of the A-B bond. In another embodiment, the mechanism of drug delivery comprises a dynamic equilibrium between B (drug moiety) bound and unbound to A (carrier moiety). The dynamic equilibrium of the carrier moiety and drug moiety to form the prodrug is:

$$A - B \leftrightarrow A + B$$

The prodrug penetrates the desired biological compartment wherein the release occurs. The drug moiety may become trapped inside the desired biological compartment. Thus, the concentration of drug moiety increases over time due to the dynamic transport by the carrier moiety which forms a labile product with the drug moiety.

A luminide may comprise a hybrid molecule of a specified biologically active agent having the optimal structure to achieve the highest therapeutic ratio and a carrier which is modified to achieve optimal bioavailability for a given drug. Thus, the physicochemical properties of a prodrug which change its bioavailability can be manipulated without altering the optimal drug structure. Luminide technology is applicable to essentially all biologically active agents. Luminides may used in any one of the following uses: antilipidemic drugs, anticholesterol drugs, aontraceptive agents, anticoagulants, anti-inflammatory agents, immuno-suppressive agents, antiarrhythmic agents, antineoplastic drugs, antihypertensive drugs, epinephrine blocking agents, cardiac inotropic drugs, antidepressant drugs, diuretics, antifungal agents, antibacterial agents, anxiolytic agents, sedatives, muscle relaxants, anticonvulsants, agents for the treatment of ulcer disease, agents for the treatment of asthma and hypersensitivity reactions, antithromboembolytic agents, agents for the treatment of muscular dystrophy,

5

10

15

20

25

30

agents to effect a therapeutic abortion, agents for the treatment of anemia, agents to improve allograft survival, agents for the treatment of disorders of purine metabolism, agents for the treatment of ischemic heart disease, agents for the treatment of opiate withdrawal, agents which activate the effects of secondary messengers, including inositol triphosphate, agents to block spinal reflexes, antiviral agents (including agents for the treatment of AIDS), pesticide applications, herbicide applications, and veterinary applications.

In an embodiment, the A moiety may be a photochromic or thermochromic molecule, e.g., the A moieties described in Table II of U.S. Patent No. 5,773,592. An exemplary A moiety is a polymethine dye or triarylmethane dye covalently bound to B. The B moiety may be any biologically active agent which covalently binds to the A moiety (e.g., as a bleaching reaction,) and wherein the bond between A and B is reversible such that the heterolytic cleavage of the covalent bond between A and B may occur inside of the desired biological compartment such as an intracellular compartment, thereby releasing B. Exemplary B moieties are shown in Table 2 herein.

Phosphonoformate (Foscarnet), an HIV-reverse transcriptase inhibitor, is an exemplary B moiety. Exemplary B moieties can also be found in Table 3 and the References at column 161 to 170 of US Patent No. 5,773,592.

Some examples of B moieties that are antihypertensive agents are tyrosine hydroxylase inhibitors such as 3,5-diiodo-4-hydrobenzoic acid; dopamine B-hydroxylase inhibitors such as mimosine and 2-mercaptoethylamine; dopa decarboxylase inhibitors such as D,L-hydrazino-α-methyldopa; and histindine decarboxylation inhibitors such as NSD 1055.

Some examples of B moieties that may be either sedatives, muscle relaxants, or anticonvulsants are neurotransmitters such as γ-aminobutyric acid; 2-oxoglutarate, aminotransferase inhibitors such as gabaculine and N-(5'-phosphopyridoxyl-4-aminobutyric acid; GABA release enhancers such as baclofen; and GABA uptake inhibitors such as trans-4-aminocrotonic acid. Some examples of B moieties that are anxoilytics are 2-oxoglutarate aminotransferase inhibitors such as gabaculine and N-(5'-phosphopyridoxyl-4-aminobutyric acid; GABA release enhancers such as baclofen; and GABA uptake inhibitors such as trans-4-aminocrotonic acid.

5

Some examples of B moieties that are anticholesterol agents are HMG-CoA reductase inhibitors such as compactin and 3-hydroxy-3methylglutarate. Some examples of B moieties that are antifungal agents are class II aldolase inhibitors such as peglycolohydroxomate and chitin synthetase inhibitors such as polyoxin D. An example of a B moiety that is an antibacterial agent is p-glycolohydroxomate, a class II aldolase inhibitor. An example of a B moiety that is an antineoplastic agent is the aspartate transcarbamylase inhibitor N-(phosphonacetyl)-L-asparatate.

An exemplary A moiety is 1,5-di-(p-N-ethyl-N-ethylaminophenyl)-1,5-bis-(p-N,N-dimethylaniline)-1,3-pentadiene, which is shown in Table 1.

Table 1: Exen	nplary A Moiety
Name	Structure
1,5-di-(p-N-ethyl-N-ethylaminophenyl)-1,5-bis-(p-N,N-dimethylaniline)-1,3-pentadiene	H_3CH_2C N $N(CH_3)_2$

Table 2: Exemp	olary B Moieties
Name	Structure
phosphonoformate	-O -P -C -O-
3,5-diiodo-4-hydrobenzoic acid	но
γ-aminobutyric acid	H ₂ NCH ₂ CH ₂ COOH
gabaculine	COOH NH ₂

Table 2: Exemp	lary B Moieties		
Name	Structure		
N-(5'-phosphopyridoxyl-4-aminobutyric acid	ОН ₂ ОР—ОН ОН ОН СН ₂ NHCH ₂ CH ₂ COOH		
baclofen	CI————————————————————————————————————		
trans-4-aminocrotonic acid	H ₂ NCH ₂ CH—CHCOOH		
compactin	COOH OH CH ₃		
3-hydroxy-3-methylglutarate	OH CH ₃ CH ₂ CCH ₂ COOH CH ₃		
P-glycolohydraoxamate	O—P—O—CH ₂ —C—N—OH		
N-(phosphonacetyl)-l-aspartate	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		
phosphonoacetate	-O—P——CH₂—CO- O-		

Table 2: Exemp	olary B Moieties	
Name	Structure	
	NH ₂ CH ₂ CHCOOH	
mimosine	ОН	
2-mercaptoethylamine	HSCH ₂ CH ₂ NH ₃ ⁺	
NSD 1055	Br——CH ₂ ONH ₂	
polyoxin D.	O OH COOH NH ₂ COCH ₂ CHCHCHCNHCH OH NH ₂ O H H OH OH OH OH	
D,L-hydrazino-α-methyldopa	HO————————————————————————————————————	

An exemplary luminide, i.e., A-B compound, is 1-phosphonoformate,1,5-di-(p-N-ethyl-N-ethylaminophenyl)-1,5-bis-(p-N,N-dimethylaniline)-1,3-pentadiene shown in Table 3.

Tab	ole 3
Name	Structure
1-phosphonoformate,1,5-di-(p-N-ethyl-N-ethylaminophenyl)-1,5-bis-(p-N,N-dimethylaniline)-1,3-pentadiene	H_3CH_2C N $N(CH_3)_2$ $N(CH_3)_2$ $N(CH_3)_2$ $N(CH_3)_2$ $N(CH_3)_2$

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than

routine experimentation, numerous equivalents to the specific procedures described herein. Such equivalents are considered to be within the scope of the present invention and are covered by the following claims. The contents of all references, issued patents, and published patent applications cited throughout this application are hereby incorporated by reference. The appropriate components, processes, and methods of those patents, applications and other documents may be selected for the present invention and embodiments thereof.

CLAIMS

What is claimed is:

1. A chemical compound having the formula A-B, where

A is a moiety capable of receiving energy; and

5

B is a biologically active agent covalently bonded to B, wherein the bond between A and B is capable of heterolytic cleavage upon reception of energy by A.

- 2. The chemical compound of claim 1, wherein A is a photochromic moiety.
- 3. The chemical compound of claim 1, wherein A is a
- thermochromic moiety.
 - 4. The chemical compound of claim 1, wherein said energy causes A to achieve an excited state; and relaxation of said excited state of A causes heterolytic cleavage of said covalent bond between A and B.
- 5. The chemical compound of claim 1, wherein said chemical compound is 1-phosphonoformate,1,5-di-(p-N-ethyl-N-ethylaminophenyl)-1,5-bis-(p-N,N-dimethylaniline)-1,3-pentadiene.
 - 6. The chemical compound of claim 1, wherein A is 1,5-di-(p-N-ethyl-N-ethylaminophenyl)-1,5-bis-(p-N,N-dimethylaniline)-1,3-pentadiene.
- 7. The chemical compound of claim 1, wherein A comprises a cationic dye20 which demonstrates photochromic behavior with electromagnetic radiation and bleaching agents.
 - 8. The chemical compound of claim 7, wherein said cationic dye is selected from the group consisting of di or triarylmethane dye, a triarylmethane lactone or a cyclic ether dye, a cationic indole, a pyronine, a phthalein, an oxazine, a thiazine, an

acridine, a phenazine, an anthocyanidin, a cationic polymethine dye, an azo or a diazopolymethine, a styryl, a cyanine, a hemicyanine, and a dialkylaminopolyene.

9. The chemical compound of claim 1, wherein B is a drug moiety which effects a therapeutic functional change by a mechanism selected from the group consisting of receptor mediated mechanisms and nonreceptor mediated mechanisms.

5

10

15

- 10. The chemical compound of claim 1, wherein B is selected from the group consisting of antilipidemic drugs, anticholesterol drugs, anticoagulants, antihypertensive drugs, cardiac inotropic drugs, antineoplastic drugs, antidepressant drugs, agents for the treatment of asthma and hypersensitivity reactions, diuretics, antifungal agents, antibacterial drugs, anxiolytic agents, sedatives, muscle relaxants, anticonvulsants, agents for the treatment of ischemic heart disease, agents which activate the effects of secondary messengers, agents to block spinal reflexes, and antiviral agents.
- 11. The chemical compound of claim 1, wherein B is selected from the group consisting of 3,5-diiodo-4-hydroxybenzoic acid, γ-aminobutyric acid, gabaculine, N-(5'-phosphopyridoxyl)-4-aminobutyric acid, baclofen, trans-4-aminocrotonic acid, compactin, 3-hydroxy-3-methylglutarate, p-glycolohydroxamate, N-(phosphonoacetyl)-L-aspartate, phosphonoacetate, mimosine, 2-mercaptoethylamine, NSD 1055, polyoxin D, D,L-hydrazino-α-methyldopa, and phosphonoformate.
- 12. The chemical compound of claim 1, wherein B is phosphonoacetate or phosphonoformate (Foscarnet).
 - 13. The chemical compound of claim 1, wherein B is an antihypertensive agent.

14. The chemical compound of claim 13, wherein B is selected from the group consisting of 3,5-diiodo-4-hydroxybenzoic acid, mimosine, D,L-hydrazino-α-methyldopa, and 2-mercaptoethylamine.

- 15. The chemical compound of claim 1, wherein B is an antiviral agent.
- 5 16. The chemical compound of claim 15, wherein B is selected from the group consisting of phosphonoacetate and phosphonoformate.
 - 17. The chemical compound of claim 1, wherein B is an anticholesterol agent.
- 18. The chemical compound of claim 17, wherein B is selected from thegroup consisting of compactin and 3-hydroxy-3-methylglutarate.
 - 19. The chemical compound of claim 1, wherein B is an anticonvulsant agent.
 - 20. The chemical compound of claim 19, wherein B is selected from the group consisting of gabaculine, N-5'-phosphopyridoxyl-4-aminobutyric acid, baclofen, trans-4-aminocrotonic acid, and γ-aminobutyric acid.
 - , 21. The chemical compound of claim 1, wherein B is an antibacterial or antifungal agent.

15

20

- 22. The chemical compound of claim 21, wherein B is selected from the group consisting of p-glycolohydroxamate and polyoxin D.
 - 23. The chemical compound of claim 1, wherein B is an antineoplastic agent.
- 24. The chemical compound of claim 23, wherein B is N-(phosphonoacetyl)-L-aspartate.
- 25. The chemical compound of claim 1, wherein said chemical compound is covalently bound to a biocompatible polymer to which an enzyme is immobilized.

26. The chemical compound of claim 25, wherein B is insulin and said enzyme is glucose oxidase.

- 27. The chemical compound of claim 25, wherein B is tissue plasminogen activator and said enzyme is xanthine oxidase.
- 28. A method for selectively delivering a biologically active agent to a biological compartment, which comprises:

5

10

15

introducing a chemical compound having the formula A-B, where A is a carrier moiety and B is a biologically active agent, into said biological compartment; and

- exposing said chemical compound to an energy sufficient to cause heterolytic cleavage of said bond between A and B, thereby releasing B from A.
 - 29. The method of claim 28, wherein A is a photochromic moiety.
 - 30. The method of claim 28, wherein A is a thermochromic moiety.
- 31. The method of claim 28, wherein said energy causes A to achieve an excited state; and relaxation of said excited state of A causes heterolytic cleavage of said covalent bond between A and B.
 - 32. The method of claim 28, wherein said chemical compound is 1-phosphonoformate,1,5-di-(p-N-ethyl-N-ethylaminophenyl)-1,5-bis-(p-N,N-dimethylaniline)-1,3-pentadiene.
- 20 33. The method of claim 28, wherein A is 1,5-di-(p-N-ethyl-N-ethylaminophenyl)-1,5-bis-(p-N,N-dimethylaniline)-1,3-pentadiene.
 - 34. The method of claim 28, wherein A comprises a cationic dye which demonstrates photochromic behavior with electromagnetic radiation and bleaching agents.

5

10

15

20

35. The method of claim 34, wherein said cationic dye is a di or triarylmethane dye, a triarylmethane lactone or a cyclic ether dye, a cationic indole, a pyronine, a phthalein, an oxazine, a thiazine, an acridine, a phenazine, an anthocyanidin, a cationic polymethine dye, an azo or a diazopolymethine, a styryl, a cyanine, a hemicyanine, or a dialkylaminopolyene.

- 36. The method of claim 28, wherein B is phosphonoformate (Foscarnet).
- 37. The method of claim 28, wherein B is selected from the group consisting of 3,5-diiodo-4-hydroxybenzoic acid, γ-aminobutyric acid, gabaculine, N-(5'-phosphopyridoxyl)-4-aminobutyric acid, baclofen, trans-4-aminocrotonic acid, compactin, 3-hydroxy-3-methylglutarate, p-glycolohydroxamate, N-(phosphonoacetyl)-L-aspartate, phosphonoacetate, mimosine, 2-mercaptoethylamine, NSD 1055, polyoxin D, D,L-hydrazino-α-methyldopa, and phosphonoformate.
- 38. The method of claim 28, wherein B is a drug molecule which effects a therapeutic functional change by a mechanism selected from the group consisting of receptor mediated mechanisms and nonreceptor mediated mechanisms.
- 39. The method of claim 28, wherein B is selected from the group consisting of antilipidemic drugs, anticholesterol drugs, anticoagulants, antihypertensive drugs, cardiac inotropic drugs, antineoplastic drugs, antidepressant drugs, agents for the treatment of asthma and hypersensitivity reactions, diuretics, antifungal agents, antibacterial drugs, anxiolytic agents, sedatives, muscle relaxants, anticonvulsants, agents for the treatment of ischemic heart disease, agents which activate the effects of secondary messengers, agents to block spinal reflexes, and antiviral agents.
 - 40. The method of claim 28, wherein B is an antihypertensive agent.

41. The method of claim 40, wherein B is selected from the group consisting of 3,5-diiodo-4-hydroxybenzoic acid, mimosine, D,L-hydrazino-α-methyldopa, and 2-mercaptoethylamine.

- 42. The method of claim 28, wherein B is an antiviral agent.
- 5 43. The method of claim 42, wherein B is selected from the group consisting of phosphonoacetate and phosphonoformate.
 - 44. The method of claim 28, wherein B is an anticholesterol agent.
 - 45. The method of claim 44, wherein B is selected from the group consisting of compactin and 3-hydroxy-3-methylglutarate.
- 10 46. The method of claim 28, wherein B is an anticonvulsant agent.
 - 47. The method of claim 46, wherein B is selected from the group consisting of gabaculine, N-5'-phosphopyridoxyl-4-aminobutyric acid, baclofen, trans-4-aminocrotonic acid, and γ-aminobutyric acid.
 - 48. The method of claim 28, wherein B is an antibacterial or antifungal agent.
- 15 49. The method of claim 48, wherein B is selected from the group consisting of p-glycolohydroxamate and polyoxin D.
 - 50. The method of claim 28, wherein B is an antineoplastic agent.
 - 51. The method of claim 50, wherein B is N-(phosphonoacetyl)-L-aspartate.
- 52. The method of claim 28, wherein said chemical compound is covalently bound to a biocompatible polymer to which an enzyme is immobilized.
 - 53. The method of claim 52, wherein B is insulin and said enzyme is glucose oxidase.
 - 54. The method of claim 52, wherein B is tissue plasminogen activator and said enzyme is xanthine oxidase.

55. The method of claim 28, wherein said energy is electromagnetic radiation.

56. The method of claim 28, wherein said energy is heat energy.

(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 20 December 2001 (20.12.2001)

PCT

(10) International Publication Number WO 01/095944 A3

(51) International Patent Classification7: A61K 41/00, 47/48

PCT/US01/18869 (21) International Application Number:

(22) International Filing Date: 12 June 2001 (12.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/211,036 12 June 2000 (12.06.2000) US

(71) Applicant and

(72) Inventor: MILLS, Randell, L. [US/US]; 493 Old Trenton Road, Cranbury, NJ 08512-5601 (US).

(74) Agents: DECONTI, Giulio, A., Jr. et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

with international search report

(88) Date of publication of the international search report: 8 August 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

IMTERNATIONAL SEARCH REPORT

Ir ational Application No
PCT/US 01/18869

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K41/00 A61K47/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, EMBASE, MEDLINE

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 09826 A (MEDIPRO SCIENCES LIMITED) 11 May 1994 (1994-05-11)	1,2,9, 10,23, 28,29, 31,38, 39,50
	page 20, line 1 - line 30 claims 1-18	
X	US 5 773 592 A (RANDER LEE MILLS) 30 June 1998 (1998-06-30) cited in the application column 7 -column 74; tables II-III claims 1-26	1-56
X	US 5 428 163 A (RANDEL L. MILLS) 27 June 1995 (1995-06-27) cited in the application column 8 -column 67/	1-56

'
χ Patent family members are listed in annex.
 "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family
Date of mailing of the international search report 19/02/2002
Authorized officer Siatou, E

INTERNATIONAL SEARCH REPORT

PCT/US 01/18869

0.40	PUI/US UI/18809		
C.(Continua Category °	ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	
Calegory	onation of document, with indication, where appropriate, or the relevant passages	nelevani to claim No.	
Х	WO 86 00527 A (DANA-FARBER CANCER INSTITUTE) 30 January 1986 (1986-01-30)	1,2,9, 10,23, 28,29, 31,38, 39,50	
x	page 7, line 1 - line 18 GOLDMACHER V S ET AL: "PHOTOACTIVATION OF	1,2,9,	
^	TOXIN CONJUGATES" BIOCONJUGATE CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 3, no. 2, 1 March 1992 (1992-03-01), pages 104-107, XP000262168 ISSN: 1043-1802 abstract	10,23, 28,29, 31,38, 39,50	
		,	
		1	

IMTERNATIONAL SEARCH REPORT

Information on patent family members

In ational Application No PCT/US 01/18869

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9409826	A	11-05-1994	US AU CA WO	5482719 A 5414194 A 2175479 A1 9409826 A2	09-01-1996 24-05-1994 11-05-1994 11-05-1994
US 5773592	A	30-06-1998	US AT AU CN DE DE EP JP JP	5428163 A 187776 T 3445489 A 1047075 A 68929117 D1 68929117 T2 0414730 A1 3025817 B2 3505574 T 8909833 A1	27-06-1995 15-01-2000 03-11-1989 21-11-1990 20-01-2000 24-08-2000 06-03-1991 27-03-2000 05-12-1991 19-10-1989
US 5428163	Α	27-06-1995	US	5773592 A	30-06-1998
WO 8600527	Α	30-01-1986	US CA DE EP JP JP WO	4625014 A 1243015 A1 3582130 D1 0185762 A1 6025071 B 61502608 T 8600527 A1	25-11-1986 11-10-1988 18-04-1991 02-07-1986 06-04-1994 13-11-1986 30-01-1986